

Residual ridge resorption, the effect on prosthodontics management of edentulous patient: an article review

Mohamed A El Maroush^a, Sarra A Benhamida^b, Ahmed A Elgendy^c, Mohamed H Elsaltani^d

^a Assistant Professor at Prosthodontics Department, Faculty of Dentistry, Tripoli University, Libya.

^b Assistant Professor at Prosthodontics Department, Faculty of Dentistry, Tripoli University, Libya.

^c Assistant Professor at Medical Physiology Department, Faculty of Medicine, Mansoura University, Egypt.

^d Assistant lecturer at Radiological Department, Faculty of Dentistry, Zantan University, Libya

Abstract:

Bone resorption is a chronic, progressive disease and irreversible process that occurs in all patients and can be classified into six different types. There are some of etiological factors that may cause resorbed ridge such as anatomical, prosthodontics, metabolic and systemic factors. The present article, were reviewed the literature concerning on possible mechanism, diagnosis, etiological factors and how to manage and prevent of resorbed ridge for edentulous patients

Key word: etiological factors, mechanism, diagnosis and management of resorbed ridge

Introduction:

The success of the complete denture relies on the fulfillment of the three basic properties which are retention, stability, and support. Mandibular dentures usually faced more difficulties in achieving these three properties than maxillary dentures, ⁽¹⁾ because, the mandible ridge has a lesser residual ridge for retention and support ⁽²⁾ and has greater resorption rate than the maxilla. According to some studies by Atwood and Tallgren show that mandibular bone resorption is four times greater than in the maxilla. ^(3,4)

Residual ridge reduction is one of the main causes of loss of denture stability and retention especially in mandibular complete dentures. Extreme resorption of the maxillary and mandibular ridges also, results in sunken appearance of cheeks, unstable and non-retentive dentures with associated pain and discomfort. Therefore, poses a clinical challenge towards the fabrication of a successful removable prosthesis. ⁽⁵⁾ Residual ridge resorption is a complex biophysical process and a common occurrence following extraction of teeth. It is the most dramatic during the first year after tooth loss followed by a slower but more progressive rate of resorption thereafter. ⁽⁶⁾ Bone resorption also is a chronic, progressive and irreversible process that occurs in all patients ⁽⁷⁾

A classification of edentulous jaw is very important as it simplify description of the residual ridge and thereby assist communication between clinicians; aid selection of the appropriate surgical prosthodontic technique; offer an objective baseline from which to evaluate and compare different treatment methods; and help in deciding on interceptive techniques to preserve the alveolar process.

A classification of the edentulous jaws has been developed based on a randomised cross-sectional study from by Cawood et al, 1988, Arising from these morphological studies of edentulous jaws they found that the basal bone does not change shape significantly, unless subjected to harmful local effects such as the overloading of ill-fitting dentures. Moreover the alveolar bone changes shape significantly in both the horizontal and vertical axes following a predictable pattern ⁽⁸⁾. Cawood and Hawell classify the residual ridge to basic six classes:

Class I - dentate.

Class II -immediately post extraction.


Class III- well-rounded ridge form, adequate in height and width.

Class IV - knife-edge ridge form, adequate in height and inadequate in width.


Class V flat ridge form, inadequate in height and width.

Class VI - depressed ridge form, with some basilar loss evident.

Moreover they found that the Pattern of bone loss varies with sites. Anterior mandible - bone loss is vertical and horizontal (from the labial aspect). Posterior mandible - bone loss is mainly vertical. Anterior maxilla - bone loss is both vertical and horizontal (from the labial aspect). Posterior maxilla - bone loss is both vertical and horizontal (from the bu]ccal aspect) ⁽³⁾



Maxillary horizontal measurements (A). Classification of resorption of maxillary alveolar ridge: anterior (B) and posterior (C). Adapted from Cawood JI, Howell RA.⁷



Modified Cawood and Howell classification of resorption. The thicker line illustrates the amount of attached mucosa, which decreases with progressive resorption. Adapted from Cawood JI, Howell RA.⁷

Etiological factors of Reduction of Residual Ridges:

There are many factors that accelerate the residual ridge resorption, Atwood postulated the main four factors which are anatomic, prosthetic, metabolic and functional factors. The anatomical factor, the shape and size of the alveolar ridge has great effect as the well-formed broad ridges show less resorption than narrow thin ridge as the force received per unit area will be less in the former. Moreover the types of the bone show great effect on the rate of bone resorption⁽⁹⁾.


The remodeling of bone is influenced by the force that applied on it which may result of Habits factor like parafunctions hapit such bruxism.⁽¹¹⁾ and misuse of prosthesis such as Intensive denture wearing, unstable occlusal conditions, Immediate denture treatment and use of improper designed denture. However Campbell observed that patients wearing complete dentures wearer presented with smaller edentulous ridges than edentulous patients with no denture treatment ⁽¹⁰⁾. Also the patients with complete dentures there is a greater degree of mandibular resorption than maxillary resorption. ^(3,4)

Metabolic and Systemic Factors has great effect on the rate of bone resorption which include Age, race, present of systematic illnesses such as osteoporosis, nutritional status especially calcium and vitamin D,⁽⁹⁾ and the amount of time the patient has been edentulous. ⁽¹¹⁾

Periodontal disease refers to the inflammatory processes that occur in the tissues surrounding the teeth in response to bacterial accumulations, or dental plaque, on the teeth. The bacterial accumulations cause an inflammatory response from the body. The chronic and progressive bacterial infection of the gums leads to alveolar bone destruction and loss of tissue attachment to the teeth. ⁽¹²⁾ Bone resorption rate are much higher In postmenopausal older women, due to inadequate of formation of new bone tissue are associated with estrogen deficiency. ⁽¹³⁾ although in those cases the rate of bone resorption can be control by treat the causes by drugs that increase bone mineral density as Bisphosphonates, RANKL inhibitors, SERMs-selective, estrogen receptor modulators, hormone replacement therapy and calcitonin. ⁽¹⁴⁾ moreover the light weight bearing exercise tends to eliminate the negative effects of bone resorption. ⁽¹⁵⁾

Cellular Mechanisms of Bone Resorption:

Bone resorption is the process by which osteoclasts break down the tissue in bones and release the minerals, resulting in a transfer of calcium from bone tissue to the blood. Osteoclasts are multi-nucleated cells (Figure2) that contain numerous mitochondria and lysosomes and it generally present on the outer layer of the bone, just beneath the periosteum. The hallmark of the resorbing surface is the appearance of scalloped erosion, called Howship's or resorption lacuna ⁽¹⁶⁾ Attachment of the osteoclast to the osteon begins the process. The osteoclast then induces an enfolding of its cell membrane and secretes collagenase and other enzymes important in the resorption process. High levels of calcium, magnesium, phosphate and products of collagen will be released into the extracellular fluid as the osteoclasts tunnel into the mineralized bone. Osteoclasts are prominent in the tissue destruction found in psoriatic arthritis and rheumatologic disorders. ⁽¹⁷⁾



Light micrograph of an osteoclast displaying typical distinguishing characteristics: a large cell with multiple nuclei and a "foamy" cytosol.

Pathogenesis of RRR:

Immediately following the extraction, any sharp edges remaining are rounded off by external osteoclastic resorption, leaving a high well rounded residual ridge. As resorption continues from the labial and lingual aspects, the crest of the ridge becomes increasingly narrow ultimately becoming knife-edged. As the process continues, the knife-edge becomes shorter and even eventually disappears, leaving a low well rounded or flat ridge. Eventually, this too resorbs, leaving a depressed ridge.⁽¹⁸⁾

The resorption phase lasts about 8–10 days, presumably the life span of the osteoclast after completion of one resorption lacuna, the osteoclast can move along the bone surface and restart resorption or undergo apoptosis.⁽¹⁹⁾

The rate of reduction in size of the residual ridge is the maximum in the first three months and then gradually tapers off.⁽²⁰⁾ The rate of residual ridge resorption differs from person to person and even at different times and sites in the same person and also affects the function of removable prostheses, which relies greatly on the quantity and the structure of jaw bones.⁽¹⁸⁾


Assessment of residual ridge resorption:

Most of the researchers have made an attempt to analyse the changes in the form of the residual alveolar ridge using some methods such as diagnostic casts as standardized measurements, lateral cephalograms and Orthopantograms radiographic.

Orthopantograms were the method chosen to assess the bone loss. This type of radiograph is subject to variations in magnification and distortion, although these problems were minimised once we were made aware of the level of magnification of our orthopantogram.⁽²¹⁾

Today CBCT is used as method to evaluate bone loss because it has shown an absence of distortion and overlapping and the dimensions it presents are compatible with the actual size Interpretation of images acquired by CBCT for evaluation of alveolar bone loss and periodontal bone defects may lead to a new approach in the evaluation of patients with periodontal disease and prove to be an excellent resource when deciding on the most appropriate therapy.⁽²²⁾

Also, All CBCT units provide axial, coronal and sagittal multi-planar reconstructed images without magnification (figure 3); CBCT displays 3D images that are necessary for the diagnosis of intra-bony defects, furcation involvements and buccal/lingual bone loss.⁽²³⁾



Tallgren ⁽²⁴⁾ found that the mean reduction in ridge height of the mandible following tooth extraction was twice that of the maxilla during the 1st year period. The ratio of mandibular to maxillary resorption increased further, to approximately ratio of 4:1 after 7 years of edentulousness ⁽²⁵⁾. The results of this study showed that the rate of resorption was almost twice more pronounced in the mandible than in the maxilla after the five-year period of complete denture wearing. That is mainly because of that the mandibular ridge is more likely to bear higher functional forces transmitted through the dentures than the maxillary ridge. Moreover the smaller surface area of the mandible and less advantageous shape of the lower basal seat. In regard to the less marked resorption of the maxillary alveolar ridge, the resistance offered by the hard palate to forces transmitted through the maxillary dentures to the denture bearing area may play an important part ⁽²⁴⁾.

Management of residual ridge resorption:

The prosthodontics management of patient with severe residual ridge resorption can be either with or without surgical intervention.

1- prosthesis without surgical intervention

As the residual ridges resorb, the tissues become unsupported and displaceable. So it's need to modify a When use of conventional impression techniques will result in a distorted impression. Therefore, the impression technique needs to be modified.

A number of modified impression techniques for resorbed mandibular ridge have been suggested by various authors such as admixed ⁽²⁶⁾, functional ⁽²⁷⁾ all green ⁽²⁸⁾, and cocktail technique ⁽²⁹⁾. All these techniques capture the primary and secondary load-bearing areas without distortion of the residual ridge. ⁽³⁰⁾ The use of these impression techniques has the following advantages: they can be easily controlled to gain maximum coverage; ⁽²⁰⁾ they can be corrected readily; ⁽³¹⁾ they can be used to accurately determine the extent of the mucobuccal reflections; and ⁽³²⁾ they can be used to direct pressure toward the load-bearing areas, specifically, the buccal shelf and the slopes of residual ridges in the mandible. ⁽³³⁾

The other technique for increases retentive and stability of complete denture is neutral zone technique. It uses with patients with highly atrophic mandible. So it of the most effective techniques is to counteract the problems of positioning the posterior teeth leading to a stable denture ⁽³⁴⁾.

Also the use of soft liners material for complete denture became popular in dentistry because they have brought many clinical advantages. These materials have the ability to help in healing of the inflamed mucosa, distribute the functional load in the support area of the prostheses and improve their adaptation and retention of complete denture ⁽³⁵⁾. Moreover, due to cushioning effect provided by soft liners, lesser amount of forces are transferred to the underlying bone during various functions as compared to one without soft liners, This leads to a reduction in residual ridge resorption.

In order to preserve the alveolar ridge and reduce the amount of stress transferred, certain general principles must be kept in mind during fabrication of complete denture. This may be achieved by having broad area of coverage under the denture base (to reduce the force per unit area) ⁽³⁶⁾. A decrease in the number of denture teeth; decrease in the buccolingual width of teeth; improved occlusal tooth design form (to decrease the amount of force required to penetrate a bolus of food) are some of the other techniques that may also be used. ⁽³⁷⁾ During tooth setup the aim should be to reduce the number of inclined planes (to minimize dislodgement of dentures and shear forces) and achieve a centralization of occlusal contacts (to increase stability of dentures and to maximize compressive load). Accurate recording of maxillomandibular relationship will ensure optimum vertical rest dimension which will decrease the frequency and duration of tooth contacts, thereby giving adequate rest to the underlying ridges. ⁽³⁸⁾

2-prosthesis with surgical intervention

In aim of achieve maximum retention and stability an implant-supported prosthesis is a reliable protocol in the management of complete edentulism⁽²¹⁾. Whereas, mandibular with two implant retained overdenture treatment is considered the 'gold standard' for the treatment of the edentulous mandible. This is based on the efficacy of this treatment modality as regards function, nutrition, and overall quality of life, balanced with patient preferences and expectations, treatment planning, prosthodontic management, and predicted costs.⁽³³⁾ However, the cost factor for such treatment over conventional dentures appears to be the only area of concern regarding its acceptability among all practitioners. With the increasing competition and marketing strategies adopted by the implant manufacturers, the cost of such implants will be sufficiently lowered for them to become affordable across the economic spectrum of patients. This will make implant supported prostheses a realistic option to rehabilitate all patients with poor ridges effectively and economically.⁽³⁹⁾

Also, the Bone grafting is a surgical procedure that replaces missing bone with material from patient's own body, an artificial, synthetic, or natural substitute. Bone grafting is possible because bone tissue has the ability to regenerate completely if provided the space into which it has to grow. As natural bone grows, it generally replaces the graft material completely, resulting in a fully integrated region of new bone.

Distraction osteogenesis (DO) is a method of generating new bone following a corticotomy or an osteotomy and gradual distraction. The method is based on the tension-stress principle proposed by Ilizarov.⁽⁴⁰⁾ The gradual bone distraction creates mechanical stimulation which induces biological responses and consequently bone regeneration. This is accomplished by a cascade of biological processes which may include differentiation of pluripotential cells, angiogenesis, osteogenesis, and bone mineralization.⁽⁴¹⁾

References:

- [1] Yadav B , Jayna M, Yada H, Suri S , Phogat S_, Madan R. Comparison of Different Final Impression Techniques for Management of Resorbed Mandibular Ridge: *Case Reports in Dentistry*, 2014;214: 6 : 25373.
- [2] Tinker A. Ageing in the United Kingdom-what does this mean for dentistry? *Br Dent J*. 2003; 194:369-372.
- [3] Atwood DA, Coy WA. Clinical Cephalometric and Densitometric Study of Reduction of Residual Ridges. *J Prosthet Dent*. 1971;26: 280-95.
- [4] Tallgren A. The Continuing Reduction of the Residual Alveolar Ridges in Complete Denture Wearers: A Mixed-Longitudinal Study Covering 25 years. *J Prosthet Dent*. 1972; 27: 120-32.
- [5] Krishna P, Mehra D, Prasad A. Prosthodontic Management Of Compromised Ridges and Situations. *INDIA, NUJHS*, 2014; 4: 2249-7110
- [6] Whitmyer C, Esposito S, Alperin S. Longitudinal Treatment of a Severely Atrophic Mandible: A Clinical Report. *J Prosthetic Dent*. 2003; 90: 116–20,
- [7] Carlsson GE. Clinical Morbidity and Sequelae of Treatment with Complete Dentures. *J Prosthet Dent*. 1998; 79: 17-23.
- [8] J. L Cawood and R. A. Howell." A classification of the edentulous jaws. *Int. J. Oral Maxillofac. Surg*. 1988; 17:232-236
- [9] Atwood D.A., 1971. "Reduction of residual ridges : A major oral disease entity". *J. Prosthet. Dent.*, 26 : 266-279
- [10] Campbell RL. A comparative Study of The Resorption of the Alveolar Ridges in Denture-Wearers and non-Denture-Wearers. *J Am Dent Assoc*. 1960;60:143-5
- [11] Carlsson GE. Responses of Jaw Bone to Pressure. *Gerodontology*. 2004;21:65–70.
- [12] Kirby A C, Meghji S, Nair S P, White P, Reddi K, Nishihara T, Nakashimi K, Willis A C, Sim R, Wilson M, Henderson B. The potent bone-resorbing mediator of *Actinobacillus*

- actinomycescomitans* is homologous to the molecular chaperone GroEL. *J Clin Invest.* 1995; 96:1185–1194.
- [13] Feng X McDonald, Jay M. Disorders of Bone Remodeling". *Annual Review of Pathology.* 2011; 01:1553-4006.
- [14] Russell G, Mueller G, Shipman C, Croucher P. "Clinical Disorders of Bone Resorption"*N. ovaritis Foundation Symposium.* 2001;232: 251–267;
- [15] Shanb AA, Youssef EF. The impact of adding weight-bearing exercise versus nonweight bearing programs to the medical treatment of elderly patients with osteoporosi . *Journal of Family and Community Medicine.* 2014;**21** (3): 176–181.
- [16] Mensah, Kofi A.; Schwarz, Edward M.; Ritchlin, Christopher . "Altered Bone Remodeling in Psoriatic Arthritis" *Current Rheumatology Reports.* 2008;**10** (4): 311–317.
- [17] Mensah, Kofi A.; Schwarz, Edward M.; Ritchlin, Christopher T. (-08-01). "Altered Bone Remodeling in Psoriatic Arthritis" *Current Rheumatology Reports* . 2008;10 (4): 311–317.
- [18] Gupta A, Tiwari B. Residual Ridge Resorption: A Review. *J India Dental Sci.* 2010; 2: 7
- [19] Teitelbaum S L. Osteoclasts: what do they do and how do they do it? *the American Journal of Pathology*,vol.170,no.2,pp. 427–435, 2007
- [20] Souza D. Oral Health Care – Prosthodontics, Periodontology, Biology, Research and Systemic Conditions
- [21] . Kreisler M, Behneke N, Behneke A, d’Hoedt B. Residual ridge resorption in the edentulous maxila in patients with implant-supported mandibular overdentures: an 8-years retrospective study. *Int J Prosthodont.* 2003;16:265-300.
- [22] Acar B, Kamburoglu K. Use of cone beam computed tomography in periodontology. *World J Radiol.* 2014. May 28; 6 (5): 139– 47.
- [23] de Faria Vasconcelos K, Evangelista KM, Rodrigues CD, Estrela C, de Sousa TO, Silva MA. Detection of periodontal bone loss using cone beam CT and intraoral radiography. *Dentomaxillofac Radiol.* 2012. January; 41 (1): 64– 9.
- [24] Lammie GA. Aging Changes and the complete lower denture. *J Prosthet Dent* 1956; 6: 450-464
- [25] TALLGREN A, *Acta Odontol Scand, J Prosthet Dent* 1967;25: 563-139
- [26] McCord JF, Tyson KW., “A conservative prosthodontic option for the treatment of edentulous patients with atrophic (flat) mandibular ridges,” *British Dental Journal*, vol. 182, no. 12, pp. 469–472, 1997.
- [27] S. Winkler, *Essentials of Complete Denture Prosthodontics*, AITBS, New Delhi, India, 2nd edition, 2009.
- [28] A. Tunkiwalla and S. Ram, “Management of mandibular poor foundation: conventional complete dentures,” *Dental Practice*, vol. 11, no. 5, pp. 34–37, 2013.
- [29] G. Praveen,. Gupta S,. Agarwal S,. Agarwal S. K, “Cocktail impression technique: a new approach to atwood's order vi mandibular ridge deformity,” *Journal of Indian Prosthodontist Society*, vol. 11, no. 1, pp. 32–35, 2011.
- [30] Carlsson GE. “Clinical morbidity and sequelae of treatment with complete dentures,” *Journal of Prosthetic Dentistry*, vol. 79, no. 1, pp. 17–23, 1998.
- [31] Whitmyer CS, Esposito J, Alperin S.“Longitudinal treatment of a severely atrophic mandible: a clinical report,” *Journal of Prosthetic Dentistry*, vol. 90, no. 2, pp. 116–120, 2003.
- [32] Wyatt CC. “The effect of prosthodontic treatment on alveolar bone loss: a review of the literature,” *The Journal of Prosthetic Dentistry*, vol. 80, no. 3, pp. 362–366, 1998.

- [33] Prithviraj D, Singh V, Kumar S, Shruti D. “Conservative prosthodontic procedures to improve mandibular denture stability in an atrophic mandibular ridge,” *Journal of Indian Prosthodontist Society*, vol. 8, no. 4, pp. 178–184, 2008.
- [34] Ohkubo C, Hanatani S, Hosoi T, Mizuno Y. Neutral zone approach for denture fabrication for a partial glossectomy patient: A clinical report. *J Prosthet Dent* 2000; 84: 390– 393. Mosby Co., 1978. 4
- [35] V. C. Petropoulos and B. Rashedi, “Current concepts and techniques in complete denture final impression procedures,” *Journal of Prosthodontics*, vol. 12, no. 4, pp. 280–287, 2003.
- [36] Rachmiel A, Leiser Y. The molecular and cellular events that take place during craniofacial distraction osteogenesis. *Plast Reconstr Surg Glob Open*. 2014;2:e98.
- [37] Manish K, Vinod K, Ravi G, Deepak M. Residual ridge resorption: a review. *Journal of Science and Technology* 2015; 1(4):124-128
- [38] Pendleton EC. Changes in the denture supporting tissue. *JADA* 1951; 42: 11-15. 32.
- [39] Ohkubo C, Hanatani S, Hosoi T, Mizuno Y. Neutral zone approach for denture fabrication for a partial glossectomy patient: A clinical report. *J Prosthet Dent* 2000; 84: 390– 393. Mosby Co., 1978.
- [40] Ilizarov GA. The tension-stress effect on the genesis and growth of tissues. Part I. The influence of stability of fixation and soft-tissue preservation. *Clin Orthop Relat Res*. 1989;238:249–81.
- [41] Derek D’Souza *ProsthodonticsIndia* Residual Ridge Resorption – Revisited Prosthodontic rehabilitation of resorbed ridges - Conventional complete dentures